



Fig. 1. Percentage of total dry weight of the diet of *Paramecium* sp. (mean \pm SD) fed on a mixture of prey items. The prey items were: diatoms, bacteria, copepods, rotifers, protozoans and other.



Fig. 2. Percentage of total dry weight of the diet of *Paramecium* sp. (mean \pm SD) fed on a mixture of prey items. The prey items were: diatoms, bacteria, copepods, rotifers, protozoans and other.

of the diet. The mean \pm SD of the percentage of total dry weight of the diet of *Paramecium* sp. fed on a mixture of prey items is shown in Figure 1. The mean \pm SD of the percentage of total dry weight of the diet of *Paramecium* sp. fed on a mixture of prey items is shown in Figure 2.

The mean \pm SD of the percentage of total dry weight of the diet of *Paramecium* sp. fed on a mixture of prey items is shown in Figure 3. The mean \pm SD of the percentage of total dry weight of the diet of *Paramecium* sp. fed on a mixture of prey items is shown in Figure 4.

The mean \pm SD of the percentage of total dry weight of the diet of *Paramecium* sp. fed on a mixture of prey items is shown in Figure 5. The mean \pm SD of the percentage of total dry weight of the diet of *Paramecium* sp. fed on a mixture of prey items is shown in Figure 6.

The mean \pm SD of the percentage of total dry weight of the diet of *Paramecium* sp. fed on a mixture of prey items is shown in Figure 7. The mean \pm SD of the percentage of total dry weight of the diet of *Paramecium* sp. fed on a mixture of prey items is shown in Figure 8.